

This document is one component of the full-length and abbreviated PPTA donor history questionnaire documents for source plasma organizations that do not use an approved test for antibodies to HIV that detect HIV-1 Group O. The full-length and abbreviated PPTA donor history questionnaire documents must be used collectively.

Source Plasma Industry Risk Poster II

Do NOT donate PLASMA, whole blood or platelets if you...

Were **born in, lived in, or travelled to** any of the following **African countries** or had **sexual contact** with anyone who was born or lived in:
Cameroon, Central African Republic, Chad, Congo, Equatorial Guinea, Gabon, Niger, Nigeria, Senegal, Togo, Zambia, Benin or Kenya

Sexual contact means any of the following (whether or not a condom or barrier device was used):

- Vaginal intercourse (**contact between penis and vagina**)
- Oral sex (**mouth or tongue on someone's vagina, penis, or anus**)
- Anal intercourse (**contact between penis and anus**).

Do NOT donate PLASMA, whole blood or platelets if you...

Ever	<ul style="list-style-type: none"> • Had HIV/AIDS (see list of symptoms below) • Had a positive test for HIV (AIDS virus) • Had a positive test for hepatitis • Had hepatitis (after your 11th birthday) • Used needles to take drugs, steroids or anything not prescribed by your doctor • Used clotting factor concentrates for a bleeding disorder • Had a transplant such as organ or bone marrow 		
Since 1977	<ul style="list-style-type: none"> • Received money, drugs or other payment for sex • (Male donors) Had sexual contact with another male, even once 		
In the last 12 months	<ul style="list-style-type: none"> • Have given money, drugs or other payment for sex • Have been treated for <ul style="list-style-type: none"> • syphilis • gonorrhoea • “Lived with” a person who has hepatitis (lived at same residence and shared kitchen and bathroom) • Had a blood transfusion or received other blood products • Received during surgery bone, tissue or skin • Had an accidental needle-stick involving exposure to blood • Had contact with someone else's blood • Had a tattoo applied • Had ear or body piercing • Have been in <table border="1" style="display: inline-table; vertical-align: middle; margin-left: 10px;"> <tr> <td style="padding: 2px;"> <ul style="list-style-type: none"> • juvenile detention • lock up • jail • prison </td> <td style="padding: 2px; vertical-align: middle; font-weight: bold;">For more than 72 hours</td> </tr> </table> <p>Had sexual contact with anyone who:</p> <ul style="list-style-type: none"> • Has HIV/AIDS (see list of symptoms below) • Has a positive test for HIV (AIDS virus) • Has hepatitis • Used needles to take drugs, steroids or anything not prescribed by their doctor • Has hemophilia or has used clotting factor concentrates • (Female donors) Had sexual contact with a male who has had sexual contact with another male, even once 	<ul style="list-style-type: none"> • juvenile detention • lock up • jail • prison 	For more than 72 hours
<ul style="list-style-type: none"> • juvenile detention • lock up • jail • prison 	For more than 72 hours		

Signs or symptoms of HIV/AIDS:

- | | |
|---|---|
| <ul style="list-style-type: none"> • Unexplained weight loss • Night sweats • Blue or purple spots in your mouth or skin • White spots or unusual sores in your mouth | <ul style="list-style-type: none"> • Swollen lymph nodes for more than one month • Fever of more than 100.5 °F for more than 10 days • Cough that won't go away • Shortness of breath • Diarrhea that won't go away |
|---|---|